

Macroscopic quantum coherence in mesoscopic ferromagnetic systems

S. P. Kou^{1,2}, J. Q. Liang^{1,2}, Y.B. Zhang^{1,2} and F. C. Pu^{1,3}

¹*Institute of Physics and Center for Condensed Matter Physics, Chinese Academy of Sciences, Beijing 100080, P. R. China*

²*Department of Physics, Shanxi University, Shanxi 030006, P. R., China*

³*Department of Physics, Guangzhou Normal College, Guangzhou 510400, P.*

In this paper we study the Macroscopic Quantum Oscillation (MQO) effect in ferromagnetic single domain magnets with a magnetic field applied along the hard anisotropy axis. The level splitting for the ground state, derived with the conventional instanton method, oscillates with the external field and is quenched at some field values. A formula for quantum tunneling at excited levels is also obtained. The existence of topological phase accounts for this kind of oscillation and the corresponding thermodynamical quantities exhibit similar interference effects which resembles to some extent the electron quantum phase interference induced by gauge potential in the Aharonov-Bohm effect and the Θ -vacuum in Yang-Mills field theory..

PACS number(s): 75.10.Jm, 03.65.Sq, 73.40.Gk, 75.30.Gw.

I. INTRODUCTION

The remarkable macroscopic quantum phenomena (MQP) of nano-magnetic particles at sufficiently low temperatures stimulated considerable research activities in recent years [1–3]. In single-domain particles, MQP represents quantum transitions of the magnetic moment $\mathbf{M}$ between easy directions in a single-domain ferromagnetic (FM) grain and of the Néel vector $\mathbf{L}$ in a quantum superposition of AFM sublattices. It has been pointed out that there is a crossover at T_B , above which the transition process is dominated by thermal hopping and the transition rate follows Arrhenius law $\Gamma = \nu_0 \exp[-U/k_B T]$ where U is the energy barrier and ν_0 is an attempt frequency, below T_B the transitions occur by quantum tunneling and a temperature independent rate $\Gamma = \nu_0 \exp[-B]$ is expected. The crossover temperature is defined as $T_B = \frac{U}{Bk_B}$. Macroscopic quantum coherence (MQC) phenomena have been studied extensively for its exotic characters far from that of classical systems. In 1992, Awschalom et al. [4] observed evidences of resonant tunneling in dense ferritin solutions in which each particle contains about 4500 Fe^{3+} spin 5/2 ions below 200mK. Recently remarkable quantization phenomena were observed in $Mn_{12}Ac$ [5] that are also understood as a kind of resonant magnetization tunneling. And resonant tunneling evidences are also shown in Fe_8 as well as CrR_6 where R stand for Mn or Ni [6,7].

It has been shown that for a wide range of magnetic systems, quantum tunneling is completely suppressed if the total spin is a half integer but is allowed for integer-spin particles which is called the "spin-parity effect" [8–10] and related to Kramers' degeneracy.. The "spin-parity effect" is a direct result of the topological phase $e^{i\phi S}$ [11] and as such does not occur in previous Lagrangian formulations of "micromagnetics", which study the statics and thermodynamics of continuous magnetization fields. In this paper we study a kind of MQC phenomena resulting from the topological phase, which we call macroscopic quantum oscillation(MQO). This kind of oscillation has been proposed in both ferromagnetic (FM) grains and antiferromagnetic molecular magnets in the presence of magnetic fields [12–14]. It has been pointed that in ferromagnetic particles with biaxial symmetry and an external field along the hard axis the tunneling rate-quenching need not be related to Kramers' degeneracy [12]. Using spin-coherent-state path integrals,

Garg first showed that the tunneling splitting is quenched [12] whenever $H = \frac{2K_1 \sqrt{1 - \frac{K_2}{K_1}}}{g\mu_B} (S - n - 1/2)$ where K_1 , K_2 are the anisotropy constants with $K_1 > K_2$, S is the total spins of the particles and n is an integer. In this paper we solve the paths from the classical action and derive a formula for the level splitting as a function of field, from which his specific field values can be read off directly. Following ref. ([14]), we calculate the tunneling dynamics of the magnetic moment of FM particles which may be measured via the static magnetization and the specific heat. Here we shall study a spin model with biaxial anisotropy in detail and point out that quantum oscillations are of the Aharonov-Bohm-type, similar to that of the θ vacuum in a Yang-Mills field but different from that in $Mn_{12}Ac$.

In Section 2 we show that there is Aharonov-Bohm-type MQO in spin systems with a uni-axial anisotropy. In section 3 we discuss a spin model with bi-axial anisotropy and find that the energy level splitting of the ground state varies with a vacuum angle Θ , which is a good realization of the Θ vacuum in condensed matter. The instanton solution is obtained through the standard procedure and the energy splitting agrees excellently with numerical results. We derive a compact formula for the level-splitting induced by tunneling which is valid for the region of low energy levels. The results of tunneling effects association with the excited states based on the LSZ (Lehmann, Symanzik

and Zimmermann) procedure of field theory are restricted in the low energy region [15–18]. Finally we study the thermodynamic properties of the tunneling effect in Section 5.

II. MQO IN MAGNETIC GRAINS WITH A EASY-AXIS ANISOTROPY

We consider the following Hamiltonian operator of a ferromagnetic particle with uniaxial anisotropy [19]

$$\mathcal{H} = K\hat{s}_z^2 - g\mu_B H\hat{s}_z \quad (1)$$

which describes uniaxial anisotropy and an magnetic field along hard axis with the anisotropy constants $K > 0$. The spin operators $\hat{s}_i$, $i = x, y, z$, obey the usual commutation relation $[\hat{s}_i, \hat{s}_j] = i\epsilon_{ijk}\hat{s}_k$ (using natural units throughout). The Hamiltonian is exactly diagonal in terms of the eigenstates $\hat{s}_z$ and we have $E_m = Km^2 - g\mu_B Hm$. Instead of using the result, we express the partition function as a spin-coherent-state path integral for large spins $S \gg 1$ [20]

$$Z = \text{Tr} e^{-\beta\mathcal{H}} = \int \mathcal{D}\{\mu(\vec{n})\} e^{-S_E} \quad (2)$$

where $\mathcal{D}\{\mu(\vec{n})\} = \prod_{k=1}^{M-1} \frac{\sin\theta_k d\phi_k d\theta_k}{2\pi}$ and

$$S_E = \int_0^{\beta\hbar} d\tau [iS\dot{\phi}(1 - \cos\theta) + KS(S+1)\cos^2\theta - g\mu_B HS\cos\theta]. \quad (3)$$

Although the method from spin-coherent-state seems redundant, our purpose in this part are to compare this simple model with the model in next part. After integrating over $\cos\theta$, we map the spin system onto a particle problem with Lagrangian

$$\mathcal{L} = \frac{m\dot{\phi}^2}{2} + i\Theta\dot{\phi} \quad (4)$$

where $m = \frac{1}{2K}$ and $\Theta = S\left(1 - \frac{g\mu_B H}{2KS}\right)$. This Lagrangian is just similar to that of eq.(3.1) in ref.([21]). The last term of $\mathcal{L}$ is the total imaginary time derivative and has no effect on the classical equation of motion, though it alters the canonical momentum $\Pi_\phi = m\dot{\phi} + i\Theta$. This magnetic model with uniaxial anisotropy in an external field resembles an electron moving on a conducting ring that is crossed by a flux Φ which shows remarkable Aharonov-Bohm effect [22]. With imaginary time $\tau = it$ variable, the periodic instanton configurations are classical solutions which minimize the Euclidean action under the boundary condition $\phi_n(\tau + \beta) = \phi_n(\tau) + 2\pi n$

$$\begin{aligned} \phi_n &= \frac{2\pi n}{\beta}\tau, \\ S_E &= s_0 n^2 + i2n\pi\Theta \end{aligned} \quad (5)$$

where n is the winding number characterizing homotopically nonequivalent class and $s_0 = \pi^2 k_B T / K$. The Euclidean functional integral of the partition function contains an additional summation over the homotopic number

$$Z = \sum_{n=-\infty}^{\infty} Z_n = \Theta_3(\pi\Theta, \exp(-s_0)) \quad (6)$$

where $\Theta_3(v, q)$ is the Jacobi theta function oscillating with Θ . In the following we shall be interested only in that part of the ground state energy which depends on the Θ

$$\begin{aligned} E_0 &= -k_B T \ln Z \text{ as } T \rightarrow 0 \\ &= -\frac{(g\mu_B H)^2}{4K} + \frac{1}{2m}\{\{\Theta\}\}^2 \end{aligned} \quad (7)$$

where $\{\{x\}\}$ is the fractional part of x to the nearest integer. This simple model shows remarkable similarity to θ vacuum of the non-Abelian gauge field - both of them are a manifestation of vacuum-vacuum tunneling and angle Θ can be made as an external variable parameter in stead of a dynamic one.

From the partition function we find the oscillation of the magnetic momentum $\frac{\partial M}{\partial H}$ with external field H_z along the hard axis

$$\begin{aligned}\frac{\partial M}{\partial H} &= -\frac{k_B T \partial^2 \ln Z}{\partial H^2} \\ &\rightarrow \frac{(g\mu_B)^2}{2K} + g\mu_B \left[\frac{1}{2} + \frac{1}{2}(-1)^{\{ \Theta \}} \right] \text{ as } k_B T m \ll 1 \\ &\rightarrow \frac{(g\mu_B)^2}{2K} - 2k_B T \left(\frac{\pi g\mu_B}{K} \right)^2 \cos(2\pi\Theta) \exp(-s_0/\hbar) \text{ as } k_B T m \gg 1.\end{aligned}\tag{8}$$

where $\{x\}$ is the integer part of x to the nearest integer. The result shows the regular intervals of magnetic field $\Delta H = \frac{2K}{g\mu_B}$. Fig.1 shows the magnetization curves at zero temperature $M - H$: curve I is magnetization curve for magnetic particles with integer spin, curve II with half integer spin and curve III is the classical one which has no oscillations. This kind of quantum behaviors is essentially a kind of "Aharonov-Bohm effect" in mesoscopic magnetic systems resulting from Berry phase. Recently quantum steps in the hysteresis loop of $M_{n12}Ac$ crystal have been observed at regular intervals of magnetic field with period $\Delta H = \frac{K}{g\mu_B}$. However the Hamiltonian of $M_{n12}Ac$ is $H = -K\hat{s}_z^2 - g\mu_B H \hat{s}_z + H'$ where H' is the perturbation term and can't communicate with other terms, such as $\hat{s}_\pm^4$ and spin-phonon interaction. Even without the term H' , this model can't be mapped onto the above model due to the opposite sign of mass, $m = \frac{-1}{2K}$. The difference between the two kinds of MQPs is obvious: in the model here MQP is essential as a kind of "Aharonov-Bohm effect" in mesoscopic magnetic systems from Berry phase with a period $\Delta H = \frac{2K}{g\mu_B}$; while in $M_{n12}Ac$ MQC is a effect of energy level-crossing and the period $\Delta H = \frac{K}{g\mu_B}$ (The perturbation term H' is the key to understand the exotic phenomenon in $M_{n12}Ac$).

III. MQO IN MAGNETIC GRAINS WITH A BIAxIAL ANISOTROPY

Let us consider the following Hamiltonian operator of a ferromagnetic particle with XOY easy plane anisotropy and x easy axis with a magnetic field applied along z axis [12]

$$\hat{H} = K_1 \hat{s}_z^2 + K_2 \hat{s}_y^2 - g\mu_B H \hat{s}_z \tag{9}$$

where $K_1 > K_2 > 0$. Starting from the coherent state representation of the time evolution operator with Hamiltonian

$$E(\theta, \phi) = K_1 \cos^2 \theta + K_2 \sin^2 \theta \sin^2 \phi - g\mu_B H \cos \theta \tag{10}$$

and integrating $\cos \theta$, we map the spin system onto a particle problem with the effective Lagrangian

$$\mathcal{L}_{eff} = \frac{1}{2} m(\phi) \dot{\phi}^2 - V(\phi) + \Theta(\phi) \dot{\phi} \tag{11}$$

where the position dependent mass $m(\phi)$ and angle $\Theta(\phi)$ are defined by

$$\begin{aligned}m(\phi) &= \frac{1}{2K_1(1 - \lambda \sin^2 \phi)}, \\ \Theta(\phi) &= S \left(1 - \frac{H}{2K_1 S (1 - \lambda \sin^2 \phi)} \right), \\ V(\phi) &= K_2 S (S + 1) \sin^2 \phi - \frac{(g\mu_B H)^2 \lambda \sin^2 \phi}{4K_1(1 - \lambda \sin^2 \phi)}\end{aligned}\tag{12}$$

respectively. Here S is the total spin of the particles and $\lambda = \frac{K_2}{K_1}$. The term $\Theta(\phi) \dot{\phi}$ has no effect on the classical motion equation and comes from the Berry phase.

A. Level Splitting of ground states

Now we derive the level splitting of a giant spin particle in the large spin limit. Instantons in field theory of $0+1$ dimensions viewed as pseudoparticles with trajectories existing in barriers, are therefore responsible for tunneling.

Since instanton trajectories interpolate between degenerate vacua and satisfy vacuum boundary conditions, the instanton method is suitable only for the calculation of tunneling parameter $\Delta\epsilon_0$ between neighboring vacua. In the following we first consider tunneling at the vacuum level (i. e. $m = 0$), which leads to the level splitting of the ground state energy, i.e. $2\Delta\epsilon_0$. Passing to imaginary time by Wick rotation $\tau = it, \beta = iT$, the amplitude for tunneling from the initial well, say that with $n = 0$ (and $\phi_i = 0$), to the neighboring well with $n = 1$ (and $\phi_f = \pi$), is obtained by

$$\langle 0, \Phi_n = \pi | e^{-\beta\mathcal{H}} | 0, \Phi_{n+1} = 0 \rangle = \mathcal{K}_E(\phi_f = \pi, \beta; \phi_i = 0, 0) \quad (13)$$

where $|0, \Phi_n\rangle$ is the perturbation wave function of the ground state at the well denoted by Φ_n . Since the classical solution ϕ_c satisfies the equation of motion $\frac{1}{2}m(\phi_c)\dot{\phi}^2 - V(\phi_c) = 0$, we have

$$\phi_c = \arccos\left[\sqrt{\frac{(1-\lambda-u)\tanh^2\omega_0\tau}{1-u-\lambda\tanh^2\omega_0\tau}}\right] \quad (14)$$

where $\omega_0 = \sqrt{4K_1K_2S(S+1)(1-u)}$ and $u = \frac{(g\mu_B H)^2}{4K_1^2S(S+1)}$ with non-vanishing topological charge and the Euclidean action is

$$S_c = \sqrt{S(S+1)} \ln \left[\frac{\sqrt{(1-u)} + \sqrt{\lambda}}{\sqrt{(1-u)} - \sqrt{\lambda}} \right] - \sqrt{S(S+1)} \sqrt{\frac{u}{1-\lambda}} \ln \left[\frac{1 + \sqrt{\xi}}{1 - \sqrt{\xi}} \right] \quad (15)$$

where $\xi = \frac{\lambda u}{(1-\lambda)(1-u)}$ and $u < (1-\lambda)$. When $H \rightarrow 0$, we have

$$\cos^2 \phi_c = \frac{1 - \tanh^2 \omega_0 \tau}{1 - \lambda \tanh^2 \omega_0 \tau} \quad (16)$$

which is exactly the vacuum instanton solution derived previously in the absence of magnetic field [2,17,18].

The prefactor can be evaluated with the stationary phase method by expanding ϕ around the instanton trajectory ϕ_c such that $\phi = \phi_c + \eta$, where η is the small fluctuation with boundary conditions $\eta(\beta) = \eta(0) = 0$. Up to the one-loop approximation we have

$$\mathcal{K}_E = e^{-S_c} I \quad (17)$$

where

$$I = \int_{\eta(0)=0}^{\eta(\beta)=0} \mathcal{D}\eta e^{-\delta S_E} \quad (18)$$

is the fluctuation functional integral with the fluctuation action

$$\delta S_E = \int_0^\beta \eta \hat{M} \eta d\tau \quad (19)$$

where

$$\hat{M} = -\frac{1}{2} \frac{d}{d\tau} m(\phi_c) \frac{d}{d\tau} + \tilde{V}(\phi_c) \quad (20)$$

with

$$\tilde{V}(\phi_c) = \frac{1}{2} [-m'(\phi_c) \ddot{\phi}_c - \frac{1}{2} m''(\phi_c) \dot{\phi}_c^2 + V''(\phi_c)] \quad (21)$$

Here $\hat{M}(\phi_c)$ is the operator of the second variation of the action and $m'(\phi_c) = \frac{\partial m(\phi)}{\partial \phi}|_{\phi=\phi_c}$. As in the usual method of evaluating the fluctuation integral I , we expand the fluctuation variable η in terms of the eigenmodes of $\hat{M}$ and set $\eta = \sum_n C_n \psi_n$, where ψ_n denotes the n -th eigenfunction of $\hat{M}$, and express the result of the integration as an inverse square root of the determinant of $\hat{M}$. In view of the time translation symmetry of the equation of motion, the functional

integral $\mathcal{K}_E$ is not well defined when expanded about the classical solutions. The translational symmetry results in zero eigenmodes of the second variation operator $\hat{M}$ of the action. This problem can be cured by the Faddeev–Popov procedure [23]. Following the procedure of refs. [24], the one instanton contribution to the propagator in the one-loop approximation is calculated as

$$\mathcal{K}_E^{(1)} = \beta \frac{4}{\pi} (1-u-\lambda)^{-\frac{1}{2}} (1-u)^{\frac{3}{2}} S^2 K_2 e^{-\omega_0 \beta/2} e^{-S_c} e^{-i\pi\Theta} \quad (22)$$

To obtain the desired result proportional to $\sinh(\beta \triangle E_0/2)$ in eq.(13), the contributions of the infinite number of instanton and anti-instanton pairs to the one instanton contribution have to be taken into account. Interactions among instantons and anti-instantons are neglected in the dilute instanton–gas approximation. Summing over all contributions of instantons and anti-instantons, the final result of the propagator is found to be

$$\begin{aligned} \mathcal{K}_E &= \lambda^{\frac{1}{4}} \left(\frac{S}{\pi}\right)^{\frac{1}{2}} (1-u)^{\frac{1}{4}} e^{-\beta\omega_0/2} \\ &\times \sum_{n,m \geq 0}^{m+n \text{ odd}} \frac{1}{m!n!} (2D)^{m+n} e^{-S_c(m+n)} e^{-i\pi\Theta(m-n)} \\ &= \lambda^{\frac{1}{4}} \left(\frac{S}{\pi}\right)^{\frac{1}{2}} (1-u)^{\frac{1}{4}} e^{-\beta\omega_0/2} \\ &\quad \sinh[\beta D \cos(\pi\Theta) e^{-S_c}] \end{aligned} \quad (23)$$

where $D = 2^3 \left\{ \frac{K_1 K_2}{(1-u-\lambda)\pi} \right\}^{\frac{1}{2}} (1-u)^{\frac{5}{4}} \lambda^{\frac{1}{4}} S^{\frac{3}{2}}$ and $\Theta = S \left(1 - \frac{g\mu_B H}{2SK_1(1-\lambda)^{1/2}} \right)$. Compared with eq.(13) the level shift is seen to be $2\triangle\epsilon_0 \cos(\pi\Theta)$ (note that this is the shift of a single level), the tunneling splitting is

$$\triangle E_0 = 2\triangle\epsilon_0 \cos(\pi\Theta) \quad (24)$$

where

$$\triangle\epsilon_0 = 8 \left\{ \frac{K_1 K_2}{(1-u-\lambda)\pi} \right\}^{\frac{1}{2}} (1-u)^{\frac{5}{4}} \lambda^{\frac{1}{4}} S^{\frac{3}{2}} e^{-S_c/\hbar}. \quad (25)$$

The tunneling splitting is quenched ($\triangle E_0 = 0$) whenever $\Theta = k + 1/2$ or $H = \frac{2\sqrt{1-\lambda}K_1}{g\mu_B} (S - k - 1/2)/S$ where k is an integer and this is the result of topological phase : there is destructive interference between any path $\phi(\tau)$ and $\theta(\tau)$ with its time reversal pair $-\phi(\tau)$ and $\pi - \theta(\tau)$. The result is in good agreement with earlier works by Garg [12]. The original Hamiltonian operator in a magnetic field along z axis has no time-reversal symmetry for $g\mu_B H \hat{s}_z \rightarrow -g\mu_B H \hat{s}_z$ as $t \rightarrow -t$ where t is time. For this reason, this quenching in magnetic field has nothing to do with Kramers' degeneracy and is different from the quenching with half-integer spins in the absence of magnetic field.

B. Level splitting of low-lying levels and the LSZ method

A formula suitable for a quantitative of the tunneling effect at excited levels is obtained with the Lehmann, Symanzik and Zimmermann (LSZ) method in field theory in which the tunneling is viewed as the transition of n bosons induced by the usual(vacuum) instanton. The idea of a tunneling transition from one side of a potential barrier to the other has recently also been linked with the LSZ reduction mechanism of a transition from the asymptotic in-state to asymptotic out-states. In the following we use the LSZ reduction procedure in a modified way in order to calculate the tunneling in the one-instanton sector for the effective potential of the spin system, including the contribution of quantum fluctuations up to the one-loop approximation.

We recall first the case of a one-dimensional harmonic oscillator described by the Hamiltonian

$$H = \frac{1}{2}p^2 + \frac{1}{2}\omega_0^2 q^2 \quad (26)$$

for mass $m = 1$. Here q and p are dynamical observable which become operators when subjected to the Heisenberg algebra of ordinary canonical quantization. The solution of the Heisenberg equation of motion, $\ddot{q} + \omega_0^2 q = 0$, then becomes

$$q(t) = \frac{1}{\sqrt{2\omega_0}} [a e^{-i\omega_0 t} + a^\dagger e^{i\omega_0 t}] \quad (27)$$

where $a, a^\dagger$ are time-independent operators defined by the initial ($t = 0$) values of q and p , i.e.

$$\begin{aligned} q(0) &= \frac{1}{\sqrt{2\omega_0}}[a + a^\dagger], \\ p(0) &= \frac{-i\omega_0}{\sqrt{2\omega_0}}[a - a^\dagger] \end{aligned} \quad (28)$$

The operators $a, a^\dagger$ can be obtained from $q(t), p(t) = \dot{q}(t)$. Thus

$$\begin{aligned} a^\dagger &= -\frac{i}{\sqrt{2\omega}}e^{-i\omega_0 t}[\dot{q}(t) + i\omega_0 q(t)] \\ &\equiv -\frac{i}{\sqrt{2\omega}}e^{-i\omega_0 t} \overleftrightarrow{\frac{\partial}{\partial t}} q(t) \end{aligned} \quad (29)$$

and a follows with complex conjugation. One should note the extra $-$ in the definition of the symbol $\overleftrightarrow{\frac{\partial}{\partial t}}$ when acting to the left. Operators of this type are well-known in the literature. Because the "harmonic oscillator" approximation is only useful of low lying levels, the results from LSZ are restricted to low lying levels.

We now consider the $(0+1)$ dimensional theory defined by the Euclidean Lagrangian eq.(12). Crucial aspects of the LSZ procedure are its asymptotic conditions which require the theory to have an interpretation in terms of observables for stationary in coming and outgoing states. We can simulate such a situation here artificially by imagining the central barrier of the potential to be extremely high and the neighboring wells " $-$ " and " $+$ " on either side to be extremely far apart. We therefore construct appropriate functions $\phi_\pm(\tau)$ which become

$$\phi_+ = \pi - \phi_c, \quad \phi_- = \phi_c \quad (30)$$

such that the interaction fields vanish in their respective asymptotic regions, i.e.

$$\lim_{\tau \rightarrow +\infty} \phi_+(\tau) = 0, \quad \lim_{\tau \rightarrow -\infty} \phi_-(\tau) = 0 \quad (31)$$

The subscripts " $-$ " and " $+$ " here denote the wells with minima at $\Phi_0 = 0$ and $\Phi_1 = \pi$ respectively. The Euclidean creation and annihilation operators $\hat{a}_\pm^\dagger$ and $\hat{a}_\pm$ which create and annihilate an effective boson in wells " $+$ " and " $-$ " respectively are related to the interaction field operators $\phi_\pm$ by

$$\begin{aligned} \hat{a}_\pm^\dagger(\tau) &:= \sqrt{\frac{2m_0}{\omega_0}}e^{-\omega_0\tau} \overleftrightarrow{\frac{\partial}{\partial \tau}} \phi_\pm(\tau), \\ \hat{a}_\pm(\tau) &:= -\sqrt{\frac{2m_0}{\omega_0}}e^{\omega_0\tau} \overleftrightarrow{\frac{\partial}{\partial \tau}} \phi_\pm(\tau) \end{aligned} \quad (32)$$

where $:=$ represents definition and $m_0 = 1/2K_1$. From the viewpoint of the LSZ method the transition amplitude between m -th degenerate eigenstates in any two neighboring wells (here for $n = 0, 1$) is viewed as the transition of m bosons induced by the instanton of eq.(14) and is related to the tunneling parameter $\Delta\epsilon_m$ by

$$A_{f,i}^m = \langle m, \Phi_1 | e^{-\beta\hat{H}} | m, \Phi_0 \rangle = e^{-\beta\epsilon_m} \sinh \beta\Delta\epsilon_m \quad (33)$$

where $| m, \Phi_n \rangle$ is the perturbation wave function of the excited level at the well denoted by Φ_n . The transition amplitude as well as the S -matrix can be related to the Green's function through the procedure known as the LSZ reduction technique. To this end we rewrite the transition amplitude as

$$A_{f,i}^m = S_{f,i}^m e^{-\beta\omega_0} \quad (34)$$

with S -matrix element

$$S_{f,i}^m = \lim_{\substack{\tau^i \rightarrow -\infty \\ \tau^f \rightarrow \infty}} \frac{1}{m!} \langle 0 | \hat{a}_+(\tau_m^f) \dots \hat{a}_+(\tau_1^f) \hat{a}_-^\dagger(\tau_1^i) \dots \hat{a}_-^\dagger(\tau_m^i) | 0 \rangle \quad (35)$$

The S -matrix element can be evaluated in terms of the Green's function G which arises in its evaluation. Thus

$$\begin{aligned}
S_{f,i}^m &= \lim_{\substack{\tau^i \rightarrow -\infty \\ \tau^f \rightarrow \infty}} \frac{1}{m!} \prod_{l=1}^m \left(-\sqrt{\frac{2m_0}{\omega_0}} e^{\omega_0 \tau_l^f} \frac{\overleftrightarrow{\partial}}{\partial \tau_l^f} \right) \left(\sqrt{\frac{2m_0}{\omega_0}} e^{-\omega_0 \tau_l^i} \frac{\overleftrightarrow{\partial}}{\partial \tau_l^i} \right) G \\
&= \frac{1}{m!} \prod_{l=1}^m \left(\frac{-2m_0}{\omega_0} \right) e^{\omega_0(\tau_l^f - \tau_l^i)} \left[\frac{\partial^2 G}{\partial \tau_l^f \partial \tau_l^i} + \omega_0 \left(\frac{\partial G}{\partial \tau_l^f} - \frac{\partial G}{\partial \tau_l^i} - \omega_0 G \right) \right]
\end{aligned} \tag{36}$$

where the $2m$ -point Green's function is defined as usual, i.e.

$$G = \langle 0 | \hat{\phi}_+(\tau_m^f) \dots \hat{\phi}_+(\tau_1^f) \hat{\phi}_-(\tau_1^i) \dots \hat{\phi}_-(\tau_m^i) | 0 \rangle \tag{37}$$

We evaluate G by inserting complete sets of states of final and initial field configurations ϕ_f, ϕ_i . Thus

$$G(\tau^f, \tau^i) = \phi_+(\tau^f) \phi_-(\tau^i) A_{f,i}^0 \tag{38}$$

which vanishes in the limit $\tau^i \rightarrow -\infty, \tau^f \rightarrow \infty$, due to eq. (31). Thus in eq. (36) the only nonvanishing contribution in these limits results from the second derivative, then

$$\frac{\partial^2 G}{\partial \tau^f \partial \tau^i} = \frac{\partial \phi_+(\tau^f)}{\partial \tau^f} \frac{\partial \phi_-(\tau^i)}{\partial \tau^i} A_{f,i}^0 \tag{39}$$

the S -matrix element for the transition of m bosons is thus

$$S_{f,i}^m = \frac{1}{m!} \prod_{l=1}^m \left\{ \left(-\frac{2m_0}{\omega_0} \right) \left[\frac{d\phi_+(\tau_l^f)}{d\tau_l^f} \frac{d\phi_-(\tau_l^i)}{d\tau_l^i} \right] \right\} A_{f,i}^0 \tag{40}$$

The transition amplitude between degenerate ground state can be calculated from the definition, eq. (26), in terms of the tunneling parameter $\Delta\epsilon_0$, i.e. $A_{f,i}^0 = \beta \Delta\epsilon_0 \cos(\Theta\pi) e^{-\beta\omega_0/2}$. Then we have

$$A_{f,i}^m = \frac{1}{m!} q^m e^{-\omega_0 m \beta} A_{f,i}^0 \tag{41}$$

where $q = \frac{8s\lambda^{1/2}(1-u)^{3/2}}{1-\lambda-u}$. The tunneling parameter at m -th excited state is seen to be

$$\Delta\epsilon_m = \frac{1}{m!} q^m \Delta\epsilon_0 \cos(\Theta\pi) \tag{42}$$

where $\Delta\epsilon_m$ is the usual overlap integral or simply the level shift due to tunneling through any one of the barriers. For $q = \frac{8s\lambda^{1/2}(1-u)^{3/2}}{1-\lambda-u} \gg 1$ the tunneling effect is much more obvious of higher excited states. The energy levels are

$$E_{m,\xi} = \epsilon_m - \xi \Delta\epsilon_m \cos(\Theta\pi) \tag{43}$$

where $\epsilon_m = \frac{(g\mu_B H)^2}{4K_1} + (m + \frac{1}{2})\omega_0$ and ξ is an integer and here can take only either of the two values “0” and “1”.

In the context of these investigations the usual terminology of MQC refers to the resonance between neighboring degenerate wells. There is an essential difficulty related to the existing theory of quantum tunneling itself in the absence of an external magnetic field. The argument of the WKB exponential of the tunneling for a ferromagnetic particle is $2\sqrt{\lambda}s$ with $\lambda = \frac{K_2}{K_1}$, K_1 and K_2 being the hard and medium axis energies. Recently, evidence for resonant tunneling were observed in $Mn_{12}Ac$, Fe_8 and CrR_6 (R equals Mn or Ni) [5–7], in which magnetic clusters have very small spins ($S = 10 - 20$). We show some data for a ferromagnetic particle in order to demonstrate the tunneling splitting in magnetic fields H with respect to the result of tunneling for half integer-spin particles in Fig.2(a) and half integer one in Fig.2(b). The result of the instanton method is plotted with the solid line and the numerical result from exact diagonalization is shown by the dotted line. The level splitting of low-lying levels can be also obtained with periodic instanton method. For the comparison see the appendices I.

IV. OBSERVATION OF MQO

In this part we discuss the possible relevance to experimental test for the topological phase interference or spin-parity effect in the single domain FM particles.

The direct way to observe MQOs is to observe the resonant tunneling frequency. Let's denote $|n >_+$ and $|n >_-$ be the eigenstates of the same energy E_n in the (separate) right and left wells, respectively. Due to quantum tunneling, the degeneracy is lifted and there emerges an energy level splitting ΔE_n . The new eigenstates are $|n >_o = \frac{1}{\sqrt{2}}(|n >_+ - |n >_-)$ with odd parity and $|n >_e = \frac{1}{\sqrt{2}}(|n >_+ + |n >_-)$ with even parity. If we choose $|\Psi(t=0) >_+ = |n >_+$ as initial state, the probability of the system being in the other well is $|C_{kn}(t)|^2 = |\langle n >_- | n >_+(t) \rangle|^2 = \sin^2[\frac{\Delta E_n t}{\hbar}]$. The resonant tunneling frequencies $\Gamma = \frac{\Delta E_n}{\hbar} = \frac{1}{n!} |\cos(\pi\Theta)| \Delta\epsilon_0 q^n$ now with magnetic field.

Since the topological phase interference can be reflected in thermodynamic quantities, it is reasonable to study the thermodynamic properties (such as the specific heat and the magnetic susceptibility). Then we discuss thermodynamic behavior of this model at very low temperature $T \sim \Delta\epsilon_0/k_B$. At such low temperature the system can be regarded as a typical two-level system

$$Z = \sum_{m=0,\xi} e^{-\beta E_{m,\xi}} = e^{-\beta\epsilon_0} \left[e^{-\beta \cos(\Theta\pi)\Delta\epsilon_0} + e^{\beta \cos(\Theta\pi)\Delta\epsilon_0} \right]. \quad (44)$$

From the partition function we can calculate the specific heat as $C_v = k_B \left[\frac{2 \cos(\Theta\pi)\Delta\epsilon_0}{k_B T} \right]^2 \text{sech}^2 \left[\frac{2 \cos(\Theta\pi)\Delta\epsilon_0}{k_B T} \right]$ which exhibits a characteristic Schottky anomaly [14] shown in Fig. 3. The height of the peak is a constant $0.64k_B$ and the correspond temperature $T_{\max}$ exhibits oscillations with a period $\frac{2K_1\sqrt{1-\lambda}}{g\mu_B}$

$$T_{\max} = \frac{1.2}{k_B} |\cos(\Theta\pi)| \Delta\epsilon_0. \quad (45)$$

The magnetization $\mathbf{M}$ and its derivative with respect to magnetic field are found to be

$$\begin{aligned} M &= \frac{(g\mu_B)^2 H}{2K_1} + \left[2 \cos(\Theta\pi) \frac{\partial \Delta\epsilon_0}{\partial H} - 2 \Delta\epsilon_0 \sin(\Theta\pi) \frac{\partial \theta}{\partial H} \right] \\ &\quad \times \tanh \left(\frac{2 \cos(\Theta\pi) \Delta\epsilon_0}{k_B T} \right) \\ &\simeq \frac{(g\mu_B)^2 H}{2K_1} - \frac{\sin(2\Theta\pi) \Delta^2 \epsilon_0}{k_B T} \frac{g\mu_B \pi}{2K_1 S \sqrt{1-\lambda}} \end{aligned} \quad (46)$$

and

$$\frac{\partial M}{\partial H} = \frac{(g\mu_B)^2}{2K_1} - \frac{2 \cos(2\Theta\pi) \Delta^2 \epsilon_0}{k_B T} \left(\frac{g\mu_B \pi}{2K_1 S \sqrt{1-\lambda}} \right)^2 \quad (47)$$

respectively. The period of oscillation in $\frac{\partial M}{\partial H}$ is $\frac{2K_1\sqrt{1-\lambda}}{g\mu_B}$.

When the temperature is higher $\Delta\epsilon_0 \ll k_B T < \omega_0$, the excited energy levels may give contribution to the partition function and we have the partition function as the following form

$$\begin{aligned} Z &= \sum_{m,\xi} \exp \left\{ -\beta \left[\frac{(g\mu_B H)^2}{4K_1} + \left(m + \frac{1}{2}\right) \omega_0 + 2 \Delta\epsilon_m \cos(\Theta\pi + \xi\pi) \right] \right\} \\ &\simeq Z_0 + 2 [\beta \Delta\epsilon_0 \cos(\Theta\pi)]^2 \exp \left[-\frac{\beta (g\mu_B H)^2}{4K_1} \right] I_0 \left[2q e^{-\beta\omega/2} \right] \end{aligned} \quad (48)$$

where $I_0[x] = \sum \frac{1}{(n!)^2} x^{2n}$ is the modified Bessel function. We define a characteristic temperature $\tilde{T}$ that is solution to the following equation $q e^{-\omega_0/2k_B \tilde{T}} \sim 1$. Only below $\tilde{T} \sim \frac{\omega_0}{2k_B \ln(q)}$ can the system be regarded as a two-level system.

From the partition function we obtain the free energy

$$\begin{aligned} F &= -\frac{1}{\beta} \ln Z \\ &\simeq -\frac{1}{\beta} \ln Z_0 - 2 [\Delta\epsilon_0 \cos(\Theta\pi)]^2 F(T, H) \end{aligned} \quad (49)$$

where $F(T, H) = \frac{I_0[2qe^{-\omega_0/2k_B T}]}{k_B T(e^{\omega_0/k_B T} - 1)}$. The magnetization $\mathbf{M}$ and its derivative with respect to magnetic field are found to be

$$M \simeq \frac{(g\mu_B)^2 H}{2K_1} - \frac{2\pi g\mu_B \sin(2\Theta\pi) \Delta^2 \epsilon_0}{2K_1 S \sqrt{1-\lambda}} F(T, H) \quad (50)$$

and

$$\frac{\partial M}{\partial H} = \frac{(g\mu_B)^2}{2K_1} - 2 \cos(2\Theta\pi) \Delta^2 \epsilon_0 \left(\frac{g\mu_B \pi}{2K_1 S \sqrt{1-\lambda}} \right)^2 F(T, H) \quad (51)$$

respectively.

Our method is only useful at low temperatures and it is not reliable above the crossover temperature T_B . Roughly speaking the picture of crossover phenomena of transition processes through the barrier can be outlined as follows. At very low temperatures (near zero) the transition is dominated by instantons or bounces which satisfy the vacuum boundary condition and are responsible for tunneling at ground state. At high temperature the transition is mainly due to the thermal activation (over barrier transition) and follows the Arrhenius law ($\sim \exp[-\frac{\Delta U}{k_B T}]$) with ΔU being the barrier height. The transition at high temperature is induced by sphelaron from the viewpoint of pseudoparticle method. The crossover occurs at certain temperature T_B above which there is only sphelaron and below which only periodic instantons contribute to the imaginary part of partition functions. Above the crossover temperature the effects of the energy levels above the barrier are more important.

V. CONCLUSION

We have discussed a spin model with biaxial anisotropy and find the energy splitting of ground state varies with an angle Θ that is a good realization of the θ vacuum structure in condensed matter. In this paper we show a kind of macroscopic quantum oscillation (MQC) phenomena in magnetic grains that is also the direct result of the topological phase and point out that some thermodynamic parameters (such as special heat C_v or dM/dH) may show oscillations in a magnetic particles with a easy-plane anisotropy, occurring in a external field along the symmetry-axis with the

period $\Delta H = \frac{K_1 \sqrt{1 - \frac{K_2}{K_1}}}{g\mu_B}$. While in a magnetic field along the medium axis, the small particles also exhibit quantum interference effects but there is no oscillation of tunneling splitting with the magnetic field. The energy levels splitting are read as the following form

$$\Delta E = \Delta E_A + \Delta E_B \cos(2S\pi) \quad (52)$$

where ΔE_A and ΔE_B have no oscillations with magnetic field [25–27].

Appendix 1: Level splitting of low-lying levels and periodic instanton method

It is by now known that the periodic instanton method has become a powerful tool for evaluation of quantum tunneling at entire region of energy. The classical solution ϕ_c satisfies the equation

$$\frac{1}{2} m(\phi_c) \dot{\phi}_c^2 - V(\phi_c) = -E_{cl} \quad (53)$$

which is obtained from by integration once and the integration constant $E_{cl} > 0$ may be viewed as the classical energy of the pseudoparticle configuration

$$\phi_c = \arcsin \sqrt{\frac{1 - k^2 \sin^2 \omega \tau}{1 - \lambda k^2 \sin^2 \omega \tau}} \quad (54)$$

where the parameters are

$$\omega = \omega_0 \sqrt{1 - n_1^{-2} n_2^2}, \quad (55)$$

$$k = \sqrt{\frac{n_1^2 - 1}{n_1^2 - n_2^2}} \quad (56)$$

and

$$\begin{aligned} n_1^2 &= 1/\left[\frac{a}{2\lambda} - \sqrt{\frac{b}{\lambda} + \frac{a^2}{4\lambda}}\right], \\ n_2^2 &= \lambda/\left[\frac{a}{2\lambda} + \sqrt{\frac{b}{\lambda} + \frac{a^2}{4\lambda}}\right] \end{aligned} \quad (57)$$

$a = -(u - \frac{4K_1\lambda E}{\omega_0^2} - 1)$ and $b = \frac{4K_1E}{\omega_0^2}$ satisfy the condition $n_1^2 > 1 > n_2^2 > 0$.

The amplitude for the transition from the left-hand well to the right-hand well at the energy E_{cl} due to instanton tunnelling can be written as

$$A_{f,i}^{E_{cl}} = \langle E_{cl}, \Phi_{n+1} | e^{-2HT} | E_{cl}, \Phi_n \rangle = e^{-2E_{cl}T} \sinh(2T\Delta E_{cl}) \quad (58)$$

and can also be evaluated with the help of the path-integral method as

$$A_{f,i}^{E_{cl}} = \int \psi_{E_{cl},n+1}^*(\phi_f) \psi_{E_{cl},n}(\phi_i) \mathcal{K}_E(\phi_f, \tau_f; \phi_i, \tau_i) d\phi_f d\phi_i \quad (59)$$

where $\psi_{E_{cl},n}(\phi_i)$, $\psi_{E_{cl},n+1}(\phi_f)$ are wave functions in the n -th, $(n+1)$ -th well respectively which extend into the domain of the barrier. ΔE_{cl} denotes the energy shift due to tunneling and is the key quantity to be calculated. Following the procedure of ref. [15–18], up to the one-loop approximation we have

$$\mathcal{K}_E = e^{-S_c} I \quad (60)$$

and the fluctuation integral can be calculated as

$$I = \frac{1}{\sqrt{2\pi}} \left[N(\tau_i) N(\tau_f) \int_{-T}^T \frac{d\tau}{N^2(\tau) m(\phi_c(\tau))} \right]^{-\frac{1}{2}} \quad (61)$$

where $N(\tau) \equiv \frac{d\phi_c}{d\tau}$ is the zero-mode of the periodic instanton solution and the Euclidean action evaluated for the periodic instanton trajectory is given by

$$\begin{aligned} S_c(\phi(T), \phi(-T), T) &= \int_{-T}^T \left(\frac{1}{2} m(\phi_c) \dot{\phi}_c^2 + V \right) d\tau \\ &= W(\phi(T), \phi(-T), T) + 2E_{cl}T + i\theta \end{aligned} \quad (62)$$

where

$$W(\phi(T), \phi(-T), E_{cl}) = \frac{\omega}{\lambda K_1} [K(k) - (1 - \lambda k^2) \Pi(\lambda k^2, k)] \quad (63)$$

and the limit $\phi(-T) \rightarrow a_1$, $\phi(T) \rightarrow a_2$ is understood. Here $K(k)$, $\Pi(\lambda k^2, k)$ are the complete elliptic integral of the first kind and the third kind, respectively. To complete the end-point integration, we choose the wave functions in the barrier as the usual WKB form

$$\begin{aligned} \psi_{E_{cl},n}(\phi_i) &= \frac{C \exp\left(-\int_{a_1}^{\phi_i} m(\phi) \dot{\phi} d\phi\right)}{\sqrt{N(\tau_i)}} \\ \psi_{E_{cl},n+1}(\phi_f) &= \frac{C \exp\left(-\int_{\phi_f}^{a_2} m(\phi) \dot{\phi} d\phi\right)}{\sqrt{N(\tau_f)}} \equiv \frac{C e^{-\Omega(\phi_f)}}{\sqrt{N(\tau_f)}} \end{aligned} \quad (64)$$

and the amplitude is written

$$\begin{aligned} A_{f,i}^{E_{cl}} &= \frac{C^2}{\sqrt{2\pi}} \int \frac{e^{-\int_{a_1}^{\phi_i} m(\phi) \dot{\phi} d\phi} e^{-\int_{\phi_f}^{a_2} m(\phi) \dot{\phi} d\phi}}{N(\tau_i)} \\ &\quad \times \left[N^2(\tau_f) \int_{-T}^T \frac{d\tau}{N^2(\tau) m(\phi_c(\tau))} \right]^{-\frac{1}{2}} e^{-S_c} d\phi_f d\phi_i \end{aligned} \quad (65)$$

The end point integration $d\phi_i$ can be evaluated as

$$\int \exp \left(- \int_{a_1}^{\phi_i} m(\phi) \dot{\phi} d\phi \right) \frac{d\phi_i}{\dot{\phi}(\tau_i)} = 2T \quad (66)$$

under the limit $\phi_i \rightarrow a_1$. We expand S_c and $\Omega = \int_{\phi_f}^{a_2} m(\phi) \dot{\phi} d\phi$ as power series of $[\phi_f - \phi(T)]$ with the limit $\phi_f \rightarrow a_2$ up to the second order for the Gaussian approximation, i.e.

$$S_c = S_c(\phi(T), \phi(-T), T) + \frac{1}{2} \frac{\partial^2 S_c}{\partial \phi_f^2} \Big|_{\phi_f=\phi(T)} (\phi_f - \phi(T))^2 + \dots \quad (67)$$

and

$$\Omega = \frac{1}{2} \frac{\partial^2 \Omega}{\partial \phi_f^2} \Big|_{\phi_f=\phi(T)} (\phi_f - \phi(T))^2 + \dots \quad (68)$$

The transition amplitude then is

$$\begin{aligned} A_{f,i}^{E_{cl}} &= \frac{2TC^2}{\sqrt{2\pi}} \int \left[\frac{1}{N^2(\tau_f) \int_{-T}^T \frac{d\tau}{N^2(\tau) m(\phi_c(\tau))}} \right]^{\frac{1}{2}} \\ &\quad \times e^{-W} e^{-2E_{cl}T} e^{-\frac{1}{2} \frac{1}{N^2(\tau_f) \int_{-T}^T \frac{d\tau}{N^2(\tau) m(\phi_c(\tau))}} (\phi_f - \phi(T))^2} d\phi_f \\ &= 2TC^2 e^{-W} e^{-2E_{cl}T} \end{aligned} \quad (69)$$

The renormalization constant C is defined and evaluated as

$$C = \left[\frac{1/2}{\int_{a_1}^{a_2} \frac{d\phi}{\sqrt{\frac{2}{m(\phi)} (E_{cl} - V)}}} \right]^{\frac{1}{2}} = \left[\frac{\omega}{4K(k')} \right]^{\frac{1}{2}} \quad (70)$$

where $k'^2 = 1 - k$. Summing over contributions from one instanton plus n instanton-anti-instanton pair, the total amplitude is given by

$$A_{f,i}^{E_{cl}} = \sum_{n=0}^{\infty} A_{f,i}^{(2n+1)} = e^{-2E_{cl}T} \sinh(2TCe^{-W}) \quad (71)$$

where the transition amplitude of one instanton plus n pairs is seen to be

$$A_{f,i}^{(2n+1)} = \int_{-T}^T d\tau_1 \int_{-T}^{\tau_1} d\tau_2 \dots \int_{-T}^{\tau_{2n}} d\tau [C]^{2n+1} e^{-(2n+1)W} e^{-2E_{cl}T} \quad (72)$$

The generalized level shift formula due to tunneling for given energy E_{cl} is obtained

$$\Delta E_m = \left[\frac{\omega}{4K(k')} \right]^{\frac{1}{2}} e^{-W} \quad (73)$$

and from which we have the energy level-splitting $2\Delta E_m \cos \pi\Theta$ where $\Theta\pi = S[1 - \frac{g\mu_B H}{2K_1 S \sqrt{1-\lambda}}]$.

- [2] E. M. Chudnovsky and L. Gunther, Phys. Rev. Lett. **60**, 661 (1988).
- [3] B. Barbara and E. M. Chudnovsky, Phys. Lett. **A145**, 205 (1990).
- [4] D. D. Awschalom, M. A. McCord, G. Grinstein, Phys. Rev. Lett. **65**, 783 (1990); D. D. Awschalom, et. al., Phys. Rev. Lett. **68**, 3029 (1992); Awschalom, et. al., Science. **258**, 414 (1992); S. Gider, D. D. Awschalom, T. Douglas, S. Mann and M. Chaparala, Science. **268**, 77 (1995).
- [5] J. R. Friedman, H. P. Sarachik, J. Tejada and R. Ziolo, Phys.. Rev. Lett. **76**, 3830 (1996); J. M. Hernandez, X. X. Zhang, F. Luis, J. Bartolome, J. Tejada and R. Ziolo, Europhys. Lett. **35**, 301 (1996); L. Thomas, F. Lioni, R. Ballou, D. Gatteschi, R. Sessoli and B. Barbara, Nature **383**, 145 (1996); J. M. Hernandez, &, Phys. Rev. **B55**, 5858 (1997).
- [6] C. Sangregorio, T. Ohm, C. Paulsen, R. Sessoli and D. Gatteschi, Phys.. Rev. Lett. **4645** (1997).
- [7] N. Vernier, G. Bellessa, T. Mallah and M. Verdaguer, Phys.. Rev. **B56**, 75 (1997).
- [8] D. Loss, D. P. Divincenzo and G. Grinstein, Phys. Rev. Lett. **69**, 3232 (1992).
- [9] J. V. Delft and C. L. Henley, Phys. Rev. Lett. **69**, 3236 (1992).
- [10] D. Loss, D. P. Divincenzo, G. Grinstein, D. D. Awschalom, and J. F. Smyth, Physica. **B189.**, 189 (1993).
- [11] E. Fradkin, *Field Theories of Condense Matter Systems* (Addison-Wesley, Reading, Ma, 1992).
- [12] A. Garg, Europhys. Lett., **22** (3), 205 (1993).
- [13] V.Yu. Golyshev and A.F. Popkov, Europhys. Lett. **29**, 327 (1992).
- [14] A. Chioloro and D. Loss, Phys. Rev. Lett. **80**, 169 (1998).
- [15] J.-G. Zhou, J.-Q. Liang, J. Burzlaff, H.J.W. Müller-Kirsten, Phys. Lett. **A223**, 142 (1996).
- [16] See for instance S. S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Harper and Row, New York, 1961).
- [17] Liang, J. Q., Müller-Kirsten, H. J. W. and Zhou, J. G. and Pu, F. C.: Phys. Lett. **A228**, 97(1997).
- [18] Liang, J. Q., Müller-Kirsten, H. J. W. and Zhou, J. G.: Z. Phys. B **102**, 525(1997).
- [19] A. Garg, Phys. Rev. **B51**, 15161 (1993).
- [20] A. Perelomov, *Generalized coherent states and their applications* (Springer, Berlin,1986).
- [21] I. V. Krive and A. S. Rozhavsky, Int. J. Mod. Phys. **B6**, 1255 (1992).
- [22] Y. Aharonov and D. Bohm, Phys. Rev. **115** , 485 (1959).
- [23] L. D. Faddeev and V. N. Popov, Phys. Lett. **B25**, 29(1967).
- [24] S. Coleman, *The Whys of Subnuclear Physics* (Plenum, New York, 1979).
- [25] M. Enz and R. Schilling, J. Phys. C: Solid State Phys. **19**, 1765 (1986).
- [26] M. Enz and R. Schilling, J. Phys. C: Solid State Phys. **19**, L711 (1986).
- [27] X.B. Wang and F.C. Pu, J. Phys.: Condens. Matter **8**, L541(1996).

Figure Captions

Fig. 1. Magnetization curves at zero temperature for a magnetic particle : curve I (solid line) is magnetization curve for magnetic particles with integer spin, curve II (dotted line) with half integer spin and curve III (dashed line) is the classical one.

Fig.2 The energy level splitting ΔE_0 of ground state for a magnetic particle with $K_1 = 1$, $\lambda = 0.2$ and (a) $S = 16$; (b) $S = 16.5$. The result of instanton method is plotted with the solid line and the numerical result is shown by the dotted line.

Fig.3 The Schottky anomaly behavior of two-level system.

Fig.1




Fig. 2(a)




Fig. 2(b)




Fig.3

